

ABUNDANCE OF YELLOW ALKALOID ‘BERBERINE’ IN THE MEDICINAL PLANT OF TALI KUNING (*Tinospora dissitiflora* Diels.) COLLECTED FROM MANOKWARI – WEST PAPUA
*(Kepadatan Alkaloid Kuning ‘Berberine’ Pada Tanaman Obat Tali Kuning (*Tinospora dissitiflora* Diels.) Asal Manokwari – Papua Barat)*

Wahyudi✉

Jurusan Kehutanan, Fakultas Kehutanan Universitas Papua Jl. Gunung Salju Amban Manokwari Papua Barat 98314, Tlp/Fax: +62986211065.

✉Penulis Korespondensi: Email: wahyudi.s.pono@gmail.com

Diterima: 12 Okt 2016 | Disetujui: 15 Nov 2016

Abstract

This research was conducting to quantify the berberine concentration. Rapid method using $^1\text{H-NMR}$ was used to determine berberine quantitative and qualitatively from the crude extract. Tali kuning was collected from Manokwari Papua Barat and powdered with hammer mill. Sonication was employed to extract using methanol at room temperature. 500 μg authentic berberine chloride were dissolved in 5.4 mL methanol-*d*4 (containing 84.4 μg anthracene). $^1\text{H-NMR}$ spectra were recorded in methanol-*d*4 (99,9%) using JEOL JNM-ECX 500. Each sample was scanned for 100 using the following parameter 0.18 Hz/point, spectral width 14400 Hz, pulse with 4.0 US, relaxation delay 2 sc. Peak areas were used for qualitative analysis and integration of each peak were employed for quantitative analysis. The results demonstrated that $^1\text{H-NMR}$ signal pattern of H-13 and H-8 recorded from Tali Kuning, and Amur corktree were well recorded, and in accordance to the berberine chloride standard. Using peak integration of H-13 and H-8, the berberine quantity in Tali kuning is 18.06 mg/g of dried powder, and 22.78 mg//gr for Amur corktree. Berberine percentage based on the weight of oven-dried-extracts was 8.34% (MC 7.54%) and 12.04% (11,54%) for Amur corktree and Tali kuning.

Keywords: Berberine, Tali kuning, Malaria, and Papuaasia

Abstract

*Penelitian ini dilakukan untuk menghitung konsentrasi berberin. Metode cepat menggunakan $^1\text{H-NMR}$ digunakan untuk menentukan jumlah berberin secara kualitatif dari ekstrak bahan mentah. Tali kuning diambil dan digiling dengan menggunakan hammar mill. Sonifikasi digunakan untuk mengekstrak methanol pada suhu kamar. Selanjutnya 500 μg berberine chlorida dilarutkan ke dalam 5.4 mL metanol-*d*4 (dengan kandungan 84.4 μg antrasin). Spektrum $^1\text{H-NMR}$ direkam dalam larutan metanol-*d*4 (99,9%) dengan menggunakan JEOL JNM-ECX 500. Setiap sampel discan dengan menggunakan parameter 0.18 Hz/point dengan lebar spektrum 14400 Hz, pulse dengan 4.0 US, relaksasi penghambatan 2 sc. Titik puncak digunakan untuk analisis kuantitatif dan integrasi dari setiap puncak. Hasil penelitian menunjukkan bahwa pola signal $^1\text{H-NMR}$ dari H-13 dan H-8 terekam dari analisis Tali Kuning dan Amur corktree juga terekam dengan baik sesuai dengan standar clorida berberin. Dengan menggunakan integrasi puncak H-13 dan H-8, jumlah berberin pada Tali kuning diperoleh 18,06 mg/g dari bubuk kering dan 22,78*

mg//gr untuk Amur corktree. Persentase kandungan berberin didasarkan pada berat ekstrak kering oven adalah 8,34% (kadar air 7,54%) dan 12,04% (11,54%) untuk Amur corktree and Tali kuning.

Kata kunci: Berberin, tali kuning, malaria, papuasia

INTRODUCTION

Several methods for quantitative and qualitative analysis of bioactive compounds in medicinal plants have been reported by several researchers, ranging from conventional methods, such as column chromatography (CC), high performance liquid chromatography (HPLC), high performance displacement liquid chromatography (HPDC) to advanced methods, like $^1\text{H-NMR}$, and liquid chromatography (LC) with electron impact or electrospray ionization mass spectrometry (EI/ESI-MS).

Advanced analytical instruments, such as $^1\text{H-NMR}$ and EI/ESI-MS etc, to determine quantities and qualities of QPAs in medicinal plants directly with extracts have been reported and employed by several researchers. Li et al. (2006; 2009) reported that rapid and simple determinations of protoberberines in the two most famous Chinese medicinal herbals, *Cortex phellodendri* (Huangbai) and *Rhizomacoptidis* (Hunglian), using $^1\text{H-NMR}$ were successfully achieved.

The $^1\text{H-NMR}$ spectra of berberine, mainly proton signals of H-13 and H-8 at ring C appeared as singlet on the specific spectrum regions (non crowded region) at δ 8.69 – 9.79 ppm, where is no interference from other signals occurred (Li et al., 2006; 2009). A suitable internal standard should be a stable compound with signal in non-crowded region of $^1\text{H-NMR}$ spectrum. For this purposes, they used anthracene ($\text{C}_{14}\text{H}_{10}$), because this compound has a stable signal at δ 8.44

ppm and its integral value is remaining constant within 48 hours.

Protoberberines are very high polar alkaloids and highly soluble in MeOH. In $^1\text{H-NMR}$ analysis, methanol- d_4 was used to avoid the interference of the phenolic hydroxyl signals (Li et al., 2006). A distinctive description of the $^1\text{H-NMR}$ for berberine and palmatine in methanol- d_4 are illustrated in Fig. 4-1, adopted from Li et al. (2006). This figure illustrates that proton signals of H-13 of berberine and palmatine appear at δ 8.69 and δ 8.79 ppm, respectively, where both resonate at non crowded-region. These clue signals could be used as key information determining protoberberine alkaloids qualification and quantification, mainly for berberine and palmatine, respectively (Li et al., 2006; 2009). The proton H-13 signals of both berberine and palmatine are well separated as singlet, and they could be used as targeted peak signals in identifying these two protoberberine alkaloids. The intensity of the signal peak is proportional to the amount of the compound (Li et al., 2006; 2009).

Li et al. (2009) made the quantitative analysis of four compounds belonging to protoberberine alkaloids in *Rhizoma coptidis*, namely berberine, palmatine, coptisine, and jatrorrhizine. The chemical structures of these compounds are shown in Fig. 4-2. On the $^1\text{H-NMR}$ analyses, the proton H-13 signals were the targeted ones for identification (Li et al., 2009). These workers suggested that these proton signals could be used as fingerprint for the identification of

protoberberine in *Rhizoma-coptidis* species as well. The $^1\text{H-NMR}$ spectra of berberine highlights that proton signal for H-13 of berberine was quite well separated from the others and at this region between δ 8.6 and 8.9 ppm, no interference by the other signals was occurred. The practitioners or traditional healers rarely use the pure compounds or single herbal drug to treat diseases, but they use a combination of several herbal drugs (Tang et al., 2009). Rapid determination of bioactive compounds in wood powder or its extract of the medicinal plants using advanced instruments, such as $^1\text{H-NMR}$, is highly desirable and recommended in identifying the herbal medicines and quantifying the main chemical constituents (Li et al., 2006; 2009). This research outlined in this paper is designed: to execute the quantitative and qualitative analysis of berberine in two medicinal plants using $^1\text{H-NMR}$, and To evaluate the medicinal value of Tali kuning (*T. dissitiflora* Diels) by comparing with well-known berberine supplier of Amur corktree (*P. amurense* Rupr).

MATERIALS AND METHODS

Two medicinal plants namely Tali kuning (*Tinosporadissitiflora* Diels) and Amur corktree (*Phelodendron Amurense* Rupr) collected from Manokwari, West Papua, and Nankoku, Kochi, Japan, respectively, were selected in this study. A quantitative and qualitative determination of berberine employed in this study was conducted using the procedure developed by Li et al. (2006; 2009). 10 mg of each powder (moisture content of 7, and 11 % respectively) were extracted with MeOH using sonication at

room temperature at duration of 30 minutes for three times. The solvent extracts were combined and dried up using vacuum evaporated to give 2.73 mg dried extract in average for Amur corktree and 1.5 mg in average for Tali kuning. Then, the dried extracts were dissolved in methanol- d_4 (0.5 ml containing 84.4 μg anthracene) and transferred into the NMR test tube directly. Three replicates were used for each medicinal plant. Berberine chloride (500 μg) dissolved in methanol- d_4 (0.5 ml containing 84.4 μg anthracene) was used as reference standard for calculating berberine content in each extract. All chemicals used in this experiment were analytical grade (MeOH, anthracene $\text{C}_{14}\text{H}_{10}$, methanol- d_4 , Wako Ltd, Osaka, Japan), and berberine chloride standard ($\text{C}_{20}\text{H}_{18}\text{NO}_4 \times \text{H}_2\text{O}$) was purchased from Tokyo Chemical Industries, Tokyo. Ultrasonic single frequency (As One Co, Osaka), NMR sample test tube 7 mm in long x 5 mm in diameter x 0.43 mm in wall thickness (Sigma-Aldrich, USA) were used, and $^1\text{H-NMR}$ was measured by JEOL JNM-ECX500 (JEOL Ltd., Japan). $^1\text{H-NMR}$ spectra were recorded in methanol- d_4 (99.9%) using JEOL JNM-ECX500 at 500 MHz. For each sample, 100 scans were conducted with the following resolution: 0.286 Hz/point; spectral width, 32768Hz; pulse width: 5.4 μs ; relaxation delay: 2s.

RESULTS AND DISCUSSION

The results of rapid determination of berberine in two medicinal plants using $^1\text{H-NMR}$ are shown in Fig. 1(A, B, and C). Figure 1 (A) demonstrates $^1\text{H-NMR}$ spectrum of the authentic berberine chloride. Whereas the $^1\text{H-NMR}$ spectra of berberine recorded for the crude extracts

of bark powders of Amur corktree and stem wood powders from Tali kuning are shown in Fig. 1 (B), and Fig. (C), respectively. These figures show that the typical berberine protons of H-13 and H-8 on the $^1\text{H-NMR}$ spectra are appeared on the downfield regions, at δ 8.69 (1H,s) and δ 9.79 (1H,s) ppm for H-13 and H-8, respectively. More importantly, these two proton signals are detected at the same chemical shifts consistently as illustrated in Fig. 4-3 (A, B, and C), respectively. These results are in well agreement and similar to those in the $^1\text{H-NMR}$ of berberine from *Cortex phellodendri* reported by Li et al. (2006).

Fig. 1 (A) also shows that the proton signal of the internal standard (anthracene) in authentic berberine chloride on the $^1\text{H-NMR}$ spectrum is detected at δ 8.44 ppm, and its integrated intensity is 1, which is equal to 84.4 μg of anthracene. Interestingly, the integrated intensities of the internal standard are completely constant at two different situations, which is 1, and appear at δ 8.44 ppm, as demonstrated in Fig. 1 (B) and (C) for Amur corktree and Tali kuning, respectively. As demonstrated in Fig. 1(A), the integrated intensity of proton H-13 of authentic berberine chloride is 1.91, which is equal to 500 μg of berberine. On the other hand, the integrated intensities of protons H-13 of berberine from the crude extracts of bark of Amur corktree shown in Fig. 1 (B) and stem wood powders of Tali kuning in Fig. 1(C) are obviously different. They are 0.84 and 0.69 for Amur corktree and Tali kuning, respectively.




Fig. 1 $^1\text{H-NMR}$ of the samples with internal standard (anthracene) in methanol- d_4 (A): Berberine chloride, (B): Extract from Amur corktree, (C): Extract from Tali kuning

From the data mentioned in the previous paragraphs, the berberine contents in the crude extracts from the bark and stem wood powders of Amur corktree and Tali kuning can be summarized in Table 1. Table 1 indicates that berberine content of the crude extracts from the bark powders of Amur corktree is 22.78 mg / g of air-dried powders, and it is higher than that (18.06 mg/g) of the crude extracts from the stem wood powders of Tali kuning. However, the percentage of berberine content in

Tali kuning based on the weight of dried extract is higher (12.04 %) than that (8.34 %) in Amur corktree (Table 4-1). It is because Tali kuning wood powder has

higher moisture content (11.15 %) and less extract content (1.50 mg) than the Amur corktree, 7.5 % and 2.73 mg, respectively.

Table 1. Quantitative determination of berberine content in the methanol extractives from bark and stem wood powders of Amur corktree and Tali kuning using integrated intensities of H-13 signals on ¹H-NMR spectra.

Medicinal plants	Moisture content (%)	Weight of dried extracts (mg) ± SD from 10 mg of air-dried wood meal	Berberine	
			Quantity (µg) ± SD in 10 mg of air-dried wood meal	Percentage based on the weight of oven-dried extract (%) ± SD
Amur corktree	7.54	2.73 ± 0.24	227.8 ± 5.7	8.34 ± 0.96
Tali kuning	11.15	1.50 ± 0.22	180.6 ± 6.4	12.04±1.90

Each value represents the value from three replicates; the following letters are standard deviation (SD)

Berberine quantity in the extracts of the stem powders of Tali kuning was comparable to that from the bark powders in Amur corktree, which is widely known for good producer of berberine (Tang et al., 2009). Based on the weight of oven-dried extracts, berberine concentration in the medicinal plants of Tali kuning (*T. dissitiflora* Diels) was higher (12.04 %) than that (8.3 %) in Amur corktree (*P. amurense* Rupr).

The quantities of berberine in the crude extracts from the two medicinal plants, Tali kuning and Amur corktree, respectively, are comparable to that (14.07 mg / g of air-dried wood meal) in *P. amurense* Rupr collected from Taiwan determined using the similar method of ¹H-NMR (Li et al., 2006). It should be considered that the whole stem of Tali kuning was used for determination of berberine content, while only the bark of Amur corktree was used for it. It means

that abundance of berberine in Tali kuning is totally considerable. Quantitative differences in berberine contents of the plants are affected by several factors, such as climate, environment, and soil composition, and the occurrence of berberine in some plants is restricted to a specific part of the plant body, but in the other plants is dispersed through the whole body with varying rates in different plant tissues (Grycova et al., 2007).

It is generally claimed that the diverse geographical origin of the plant make the content of alkaloids quite different one to another (Li et al. 2006; 2009). Therefore, a rapid and simple determination of bioactive compounds in the plants using ¹H-NMR is necessary to estimate the medicinal values of it. These results support the arguments that ¹H-NMR is an effective tool superior to HPLC in detecting and determining the chemical

constituents in herbal medicine, and has been utilized for quality control of commercial traditional Chinese one.

REFERENCES

- Li CY, HJ Lu, CH Lin, TS Wu. 2006. A Rapid and Simple Determination of Protoberberine Alkaloids in *Cortex phellodendri* by 1H-NMR and Its application for Quality Control of Commercial Traditional Chinese Medicine Prescriptions. Journal of Pharmacheutical and Biomedical Analysis 40: 173-178.
- Li CY, SI Tsai, AG Damu, TS Wu. 2009. A Rapid and Simple Determination of Protoberberine Alkaloids in *Rhizoma coptidis* by 1H-NMR and Its application for Quality Control of Commercial Prescriptions. Journal of Pharmacheutical and Biomedical Analysis 49: 1272-1276.
- Wahyudi, Y Ohtani, H Ichiura. 2011. Berberine in the Medical plant of Tali Kuning (*Tinosporadissitiflora* Diels). Journal of Wood Science 2 (2):100-104.